EXBIO Antibodies

PRODUCT DATA SHEET

1A-308-T025

Monoclonal Antibody to CD27 Allophycocyanin (APC) conjugated (25 tests)

Clone: LT27

Isotype: Mouse IgG2a

Specificity: The antibody LT27 reacts with CD27 (T14), a 50-55 kDa type I transmembrane glycoprotein (member of the TNF-receptor superfamily) expressed on medullary thymocytes, peripheral T lymphocytes, some B lymphocytes and NK cells.
HLDA V; WS Code T T-CD27.01

Immunogen: Human peripheral blood lymphocytes

Species Reactivity: Human

Preparation: The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.

Storage Buffer: The reagent is provided in phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA) as a stabilizing agent.

Storage / Stability: Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.
Do not use after expiration date stamped on vial label.
Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.

Usage: The reagent is designed for Flow Cytometry analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10^6 cells in a suspension.
The content of a vial (0.5 ml) is sufficient for 25 tests.

Expiration: See vial label

Lot Number: See vial label

Background: CD27 is a transmembrane 55 kDa protein of the nerve growth factor-receptor family, expressed as a disulfide-linked homodimer on mature thymocytes, peripheral blood T cells and a subpopulation of B cells. Activation of T cells via TCR-CD3 complex results in upregulation of CD27 expression on the plasma membrane as well as in the release of its soluble 28-32 kDa form, sCD27, detected in the plasma, urine or spinal fluid. This sCD27 is an important prognostic marker of acute and chronic B cell malignancies. RgpA, a cystein proteinase, although activating T cells through the protease-activated receptors (PARs), degradates CD27 and counteracts T cell activation mediated by CD27 and its ligand CD70.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 42 Vestec u Prahy | Czech Republic
Tel: +420 261 090 664 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz

References:

*van Oers MH, Pals ST, Evers LM, van der Schoot CE, Koopman G, Bonfrer JM, Hintzen RQ, von dem Borne AE, van Lier RA: Expression and release of CD27 in human B-cell malignancies. Blood. 1993 Dec 1;82(11):3430-6.
*Molica S, Vitelli G, Levato D, Crispino G, Dell'Olio M, Dattilo A, Matera R, Gandolfo GM, Musto P. CD27 in B-cell chronic lymphocytic leukemia. Cellular expression, serum release and correlation with other soluble molecules belonging to nerve growth factor receptors (NGFr) superfamily. Haematologica. 1998 May;83(5):398-402.
*Kara IO, Sahin B, Gunesacar R: Expression of soluble CD27 and interleukins-8 and -10 in B-cell chronic lymphocytic leukemia: correlation with disease stage and prognosis. Adv Ther. 2007 Jan-Feb;24(1):29-40.
*Yun LW, Decarlo AA, Hunter N: Blockade of protease-activated receptors on T cells correlates with altered proteolysis of CD27 by gingipains of Porphyromonas gingivalis. Clin Exp Immunol. 2007 Nov;150(2):217-29.
*Leukocyte Typing V., Schlossman S. et al. (Eds.), Oxford University Press (1995).

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 42 Vestec u Prahy | Czech Republic
Tel: +420 261 090 664 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz