EXBIO Antibodies

PRODUCT DATA SHEET

1F-306-T100

Monoclonal Antibody to CD21 Fluorescein (FITC) conjugated (100 tests)

Clone: LT21

Isotype: Mouse IgG1

Specificity: The antibody LT21 reacts with CD21 (CR2), a 145 kDa transmembrane glycoprotein (complement C3d receptor - C3dR) expressed on B lymphocytes, follicular dendritic cells, some epithelial cells and a subsets of T lymphocytes. It is not expressed on immature B cells.
HLDA VI; WS Code B CD21.1

Immunogen: IM9 human B-lymphoblastoid cell line

Species Reactivity: Human, Porcine, Bovine, Canine (Dog)

Preparation: The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.

Storage Buffer: The reagent is provided in phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA) as a stabilizing agent.

Storage / Stability: Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.
Do not use after expiration date stamped on vial label.
Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.

Usage: The reagent is designed for Flow Cytometry analysis of human blood cells using 20 µl reagent / 100 µl of whole blood or 10^6 cells in a suspension.
The content of a vial (2 ml) is sufficient for 100 tests.

Expiration: See vial label

Lot Number: See vial label

Background: CD21 (complement receptor 2, CR2) binds C3 complement fragments, especially its breakdown fragments, which remain covalently attached to complement activating surfaces or antigen. CD21 has important roles in uptake and retention of immunocomplexes, survival of memory B cells and in development and maintenance of the humoral response to T-dependent antigens. CD21 also serves as a key receptor for Epstein-Barr virus binding and is involved in targeting prions to folicular dendritic cells and expediting neuroinvasion following peripheral exposure to prions. A soluble form of the CD21 (sCD21) is shed from the lymphocyte surface and retains its ability to bind respective ligands.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 42 Vestec u Prahy | Czech Republic
Tel: +420 261 090 664 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz

References:

*Roozendaal R, Carroll MC: Complement receptors CD21 and CD35 in humoral immunity. Immunol Rev. 2007 Oct;219:157-66.
*Twohig J, Kulik L, Haluszczak C, Reuter J, Rossbach A, Bull M, Holers VM, Marchbank KJ: Defective B cell ontogeny and immune response in human complement receptor 2 (CR2, CD21) transgenic mice is partially recovered in the absence of C3. Mol Immunol. 2007 Jul;44(13):3434-44.
*Kasprzak A, Spachacz R, Wachowiak J, Stefanska K, Zabel M: Epstein-Barr virus (EBV) infection in B-cell non-Hodgkin's lymphomas in children: virus latency and its correlation with CD21 and CD23 molecules. Folia Histochem Cytobiol. 2007;45(3):169-79.
*Zabel MD, Heikenwalder M, Prinz M, Arrighi I, Schwarz P, Kranich J, von Teichman A, Haas KM, Zeller N, Tedder TF, Weis JH, Aguzzi A: Stromal complement receptor CD21/35 facilitates lymphoid prion colonization and pathogenesis. J Immunol. 2007 Nov 1;179(9):6144-52.
*Singh A, Blank M, Shoenfeld Y, Illges H: Antiphospholipid syndrome patients display reduced titers of soluble CD21 in their sera irrespective of circulating anti-beta2-glycoprotein-I autoantibodies. Rheumatol Int. 2008 Jan 3
*Leukocyte Typing VI., Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).
*Faldyna M, Samankova P, Leva L, Cerny J, Oujezdska J, Rehakova Z, Sinkora J: Cross-reactive anti-human monoclonal antibodies as a tool for B-cell identification in dogs and pigs. Vet Immunol Immunopathol. 2007 Sep 15;119(1-2):56-62.
*Filatov AV, Krotov GI, Zgoda VG, Volkov Y: Fluorescent immunoprecipitation analysis of cell surface proteins: a methodology compatible with mass-spectrometry. J Immunol Methods. 2007 Jan 30;319(1-2):21-33.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 42 Vestec u Prahy | Czech Republic
Tel: +420 261 090 664 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz