

Technical Data Sheet


Purified NA/LE Non-Cytolytic Mouse CTLA-4 - IgG Fusion Protein

Product Information

Material Number:	552133
Size:	0.5 mg
Concentration:	0.5 mg/ml
Storage Buffer:	No azide/low endotoxin: Aqueous buffered solution containing no preservative, 0.2µm filtered. Endotoxin level is ≤0.01 ng/µg of protein.

Description

The Non-Cytolytic Mouse CTLA-4 - IgG Fusion Protein is composed of the extracellular domain of mouse CTLA-4 (CD152) fused to a mutant Fc region of mouse IgG2a, which is unable to bind to the C1q complement or to Fc receptors. This chimeric protein binds to CTLA-4's ligands B7-1 (CD80) and B7-2 (CD86) on mouse antigen-presenting cells and blocks their binding to both CTLA-4 and CD28. CTLA-4 (also known as CD152) is a cell-surface Ig-superfamily glycoprotein closely related to the CD28 costimulatory receptor. CTLA-4 is expressed on activated T lymphocytes 2-3 days after stimulation through the T-cell receptor. Whereas CD28 delivers a costimulatory signal required for T-cell activation, CTLA-4 is a negative regulator of cell-mediated immune responses. CTLA-4 IgG Fusion Protein has been shown to prevent allograft rejection and induce donor-specific tolerance, inhibit *in vitro* responses of splenocytes to ConA; inhibit the spontaneous *in vitro* and *in vivo* lymphoproliferation in CTLA-4-deficient mice; and limit superantigen-induced T-cell proliferation, anergy, and secretion of IL-2, IFN-γ, and IL-4, but not TNF-α or IL-10.



Inhibition of ConA-induced proliferation of splenic T lymphocytes by Non-Cytolytic Mouse CTLA-4 - IgG Fusion Protein. C57BL/6 splenocytes were cultured for three days, either with no stimulation or with 4 µg ConA per 10⁶ cells, as indicated. The indicated concentrations of either Non-Cytolytic Mouse CTLA-4 - IgG Fusion Protein or a control fusion protein were added to ConA-stimulated cells. Cell proliferation was quantitated by the MTT fluorometric assay. The CTLA-4 fusion protein inhibited ConA-induced cell proliferation in a dose-dependent manner, whereas the control fusion protein had little effect.

Preparation and Storage

This preparation contains no preservatives, thus it should be handled under aseptic conditions.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Avoid multiple freeze-thaws of product.

The fusion protein solution should be stored at -20°C until the vial is opened. Thawed aliquots may be stored for at least 1 week at 4°C.

Application Notes

Application

Blocking	Routinely Tested
----------	------------------

Recommended Assay Procedure:

This fusion protein has been tested by LAL assay for endotoxin level, by SDS-PAGE to assure purity, and by an *in vitro* T-cell proliferation assay to assure blocking activity.

Product Notices

- Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

References

- Carreno BM, Collins M. The B7 family of ligands and its receptors: New pathways for costimulation and inhibition of immune responses. *Annu Rev Immunol.* 2002; 20:29-53.(Biology)
- Gonzalo JA, Delaney T, Corcoran J, Goodearl A, Gutierrez-Ramos JC, Coyle AJ. Cutting edge: the related molecules CD28 and inducible costimulator deliver both unique and complementary signals required for optimal T cell activation. *J Immunol.* 2001; 166(1):1-5.(Biology)
- Lenschow DJ, Walunas TL, Bluestone JA. CD28/B7 system of T cell costimulation. *Annu Rev Immunol.* 1996; 14:233-258.(Biology)
- Steurer W, Nickerson PW, Steele AW, Steiger J, Zheng XX, Strom TB. Ex vivo coating of islet cell allografts with murine CTLA4/Fc promotes graft tolerance. *J Immunol.* 1995; 155(3):1165-1174.(Biology)
- Tivol EA, Boyd SD, McKeon S, et al. CTLA4lg prevents lymphoproliferation and fatal multiorgan tissue destruction in CTLA-4-deficient mice. *J Immunol.* 1997; 158(11):5091-5094.(Biology)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD

