


Technical Data Sheet

Purified Mouse Anti-nNOS/NOS Type 1**Product Information**


Material Number:	611852
Size:	50 µg
Concentration:	250 µg/ml
Clone:	52/nNOS/NOS Type I
Immunogen:	Rat nNOS/NOS Type I aa. 144-262
Isotype:	Mouse IgG1
Reactivity:	QC Testing: Rat
Target MW:	155 kDa
Storage Buffer:	Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium azide.

Description

Nitric oxide synthase (NOS) is a cell-type specific enzyme which catalyzes the synthesis of nitric oxide (NO). NO is a short-lived radical which transmits cellular signals involved in vasorelaxation, neurotransmission, and cytotoxicity. In neurons and endothelial cells, constitutive NOS (cNOS) is activated by agonists that increase intracellular Ca²⁺ concentrations and enhance calmodulin binding. Neuronal NOS (nNOS or bNOS) and endothelial NOS (eNOS) have recognition sites for NADPH, FAD, FMN, and calmodulin and are regulated in a similar manner. However, both have been shown to be distinct gene products of about 155kDa and 140kDa, respectively, and the human forms show 52% amino acid identity to each other. nNOS and the 130 kDa inducible macrophage NOS (iNOS) share 51% amino acid homology. nNOS (neuronal NOS) is found in the cytoplasm of cell types such as neurons, skeletal muscle fibers, and lung epithelium. It is slightly larger than the other isoforms due to an N-terminal extension containing a PDZ domain through which it binds PSD-95 (post-synaptic density protein) and localizes to the cell membrane of synapses.



Western blot analysis of nNOS/NOS Type I on rat pituitary lysate. Lane 1: 1:2500, lane 2: 1:5000, lane 3: 1:10000 dilution of nNOS/NOS Type I.



GC

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Store undiluted at -20°C.

Application Notes**Application**

Western blot	Routinely Tested
Immunofluorescence	Tested During Development

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

BD Biosciences

bdbiosciences.com

United States 877.232.8995 Canada 888.259.0187 Europe 32.53.720.550 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

- Bredt DS, Hwang PM, Glatt CE, Lowenstein C, Reed RR, Snyder SH. Cloned and expressed nitric oxide synthase structurally resembles cytochrome P-450 reductase. *Nature*. 1991; 351(6329):714.(Biology)
- Nathan C. Nitric oxide as a secretory product of mammalian cells. *FASEB J*. 1992; 6(12):3051-3064.(Biology)
- Tochio H, Mok YK, Zhang Q, Kan HM, Bredt DS, Zhang M. Formation of nNOS/PSD-95 PDZ dimer requires a preformed beta-finger structure from the nNOS PDZ domain. *J Mol Biol*. 2000; 303(3):359-370.(Biology)