

Dvl-1 siRNA (bovine): sc-270669

BACKGROUND

Mammalian homologs of the *Drosophila* dishevelled (Dsh) gene have been identified, including Dvl-1, Dvl-2 and Dvl-3. The mammalian dishevelled proteins contain three homologous domains, two of which are unrelated to any other known protein. The third region is homologous to the discs-large homology domain of *Drosophila* discs-large-1, a tumor suppressor protein. Like their *Drosophila* counterpart, the dishevelled proteins are thought to be involved in embryogenesis. Overexpression of Dvl-1 has been shown to inhibit the phosphorylation of Tau by GSK-3 β . This finding may prove to be important in Alzheimer's studies, which have shown that tau is hyperphosphorylated. In *Drosophila*, Dsh is a component of the frizzled signaling pathway. Both mammalian dishevelled and frizzled proteins are components of the Wnt signalling pathway.

REFERENCES

1. Sussman, D.J., Klingensmith, J., Salinas, P., Adams, P.S., Nusse, R. and Perrimon, N. 1994. Isolation and characterization of a mouse homolog of the *Drosophila* segment polarity gene dishevelled. *Dev. Biol.* 166: 73-86.
2. Krasnow, R.E., Wong, L.L. and Adler, P.N. 1995. Dishevelled is a component of the frizzled signaling pathway in *Drosophila*. *Development* 121: 4095-4102.
3. Yang-Snyder, J., Miller, J.R., Brown, J.D., Lai, C.J. and Moon, R.T. 1996. A frizzled homolog functions in a vertebrate Wnt signaling pathway. *Curr. Biol.* 6: 1302-1306.
4. Pizzuti, A., Novelli, G., Mari, A., Ratti, A., Colosimo, A., Amati, F., Penso, D., Sangiuolo, F., Calabrese, G., Palka, G., Silani, V., Gennarelli, M., Mingarelli, R., Scarlato, G., Scambler, P. and Dallapiccola, B. 1996. Human homologue sequences to the *Drosophila* dishevelled segment-polarity are deleted in the DiGeorge syndrome. *Am. J. Hum. Genet.* 58: 722-729.
5. Tsang, M., Lijam, N., Yang, Y., Beier, D.R., Wynshaw-Boris, A. and Sussman, D.J. 1996. Isolation and characterization of mouse dishevelled-3. *Dev. Dyn.* 207: 253-262.
6. Pizzuti, A., Amati, F., Calabrese, G., Mari, A., Colosimo, A., Silani, V., Giardino, L., Ratti, A., Penso, D., Calza, L., Palka, G., Scarlato, G., Novelli, G. and Dallapiccola, B. 1996. cDNA characterization and chromosomal mapping of two human homologues of the *Drosophila* dishevelled polarity gene. *Hum. Mol. Genet.* 5: 953-958.
7. Semenov, M.V. and Snyder, M. 1997. Human dishevelled genes constitute a DHR-containing multigene family. *Genomics* 42: 302-310.
8. Wagner, U., Brownlees, J., Irving, N.G., Lucas, F.R., Salinas, P.C. and Miller, C.C. 1997. Overexpression of the mouse dishevelled-1 protein inhibits GSK-3 β -mediated phosphorylation of Tau in transfected mammalian cells. *FEBS Lett.* 411: 369-372.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic Locus: DVL1 (bovine) mapping to 16.

PRODUCT

Dvl-1 siRNA (bovine) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Dvl-1 shRNA Plasmid (bovine): sc-270669-SH and Dvl-1 shRNA (bovine) Lentiviral Particles: sc-270669-V as alternate gene silencing products.

For independent verification of Dvl-1 (bovine) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-270669A and sc-270669B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Dvl-1 siRNA (bovine) is recommended for the inhibition of Dvl-1 expression in bovine cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Dvl-1 gene expression knockdown using RT-PCR Primer: Dvl-1 (bovine)-PR: sc-270669-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.