

Mouse anti-CD19, clone LE-CD19 (Monoclonal)

Clone no. LE-CD19

MONOSAN

Product name	Mouse anti-CD19, clone LE-CD19 (Monoclonal)
Host	Mouse
Applications	WB,IHC-P,IP,FC,ELISA
Species reactivity	human
Conjugate	Purified
Immunogen	CD19 peptide CGPDPAWGGGGRMGWSTR (C-terminus) coupled to KLH.
Isotype	IgG1
Clonality	Monoclonal
Clone number	LE-CD19
Size	0.2 mg
Concentration	1.0 mg/ml
Format	-
Storage buffer	PBS with 0.09% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-CD19, clone LE-CD19 (Monoclonal)

Clone no. LE-CD19

MONOSAN

Additional info

Mouse anti Human CD19 antibody, clone LE-CD19 recognizes an epitope within the C-terminal cytoplasmic tail sequence of human CD19, a single pass type I transmembrane glycoprotein containing two C2 type Ig-like domains in the N-terminal extracellular region and four potential phosphorylation sites for tyrosine together with a single serine in the cytoplasmic region. Human CD19 is expressed on virtually all cells of the B-cell lineage with the exception of plasma cells and plays a regulatory role in B-cell differentiation and proliferation. B-cells are essential for antibody production and mutations in the CD19 gene can lead to an immunodeficiency syndrome, CIVD3 characterized by hypogammaglobulinemia leading to recurrent infections and the inability to mount an antibody mediated response to immune insult. Although immunoglobulin production is impaired B-cell precursors appear in normal numbers together with some reduction in more mature B-cell forms (van Zelm et al. 2006). B-cells have also been implicated in the progression and pathogenesis of multiple sclerosis and are common components of both active and chronic MS lesions and well as the CSF (Ritchie et al. 2004) Mouse anti Human CD19 antibody, clone LE-CD19 has been successfully employed for the immunohistochemical demonstration of CD19 in formalin fixed, paraffin embedded tissues (Streeck, H. et al. 2011) and for the detection of CD19 in cell lysates by Western blotting.

References

1. -
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES