

MEF Feeder Cells (Puromycin-resistant)

CATALOG NUMBER: CBA-312

STORAGE: Liquid nitrogen

Note: For best results begin culture of cells immediately upon receipt. If this is not possible, store at -80°C until first culture. Store subsequent cultured cells long term in liquid nitrogen.

QUANTITY & CONCENTRATION: 1 mL, 5 x 10⁶ cells/mL in 90% FBS, 10% DMSO

Background

Embryonic stem (ES) cells have been derived from the inner cell masses (ICM) of blastocysts in many species. They are capable of unlimited, undifferentiated proliferation on feeder cell layers and remain karyotypically normal and phenotypically stable. In addition, ES cells have the ability to differentiate into a wide variety of cell types *in vitro* and *in vivo*. In mES cell culture, the feeder layer can be replaced by the addition of LIF in the growth medium. However, LIF does not have the same effect on hES cell culture as mES. Therefore, both the derivation and maintenance of hES cells require the use of feeder cells.

Application

Puromycin-resistant Embryonic Fibroblast (MEF) feeder cells are used for the maintenance of stable Puromycin-resistant mouse or human ES cells in the undifferentiated state. The cells must be mitotically inactivated prior to the addition of ES cells, such as treatment with mitomycin C (2-4 hr, 10 µg/mL).

Quality Control

Puromycin-resistant Embryonic Fibroblast (MEF) feeder cells are cultured in medium containing 2 µg/mL Puromycin. This cryovial contains at least 5.0 × 10⁶ MEF feeder cells as determined by morphology, trypan-blue dye exclusion, and viable cell count. The MEF feeder cells are tested free of microbial contamination.

Medium

1. Culture Medium: D-MEM (high glucose), 10% fetal bovine serum (FBS), 2 µg/mL Puromycin, 0.1 mM MEM Non-Essential Amino Acids (NEAA), 2 mM L-glutamine, 1% Pen-Strep (optional)
2. Freeze Medium: 90% FBS, 10% DMSO

Methods

I. Establishing MEF Feeder Cell Cultures from Frozen Cells

1. Place 10 ml of complete DMEM growth medium in a 50-ml conical tube. Thaw the frozen cryovial of cells within 1–2 minutes by gentle agitation in a 37°C water bath. Decontaminate the cryovial by wiping the surface of the vial with 70% (v/v) ethanol.
2. Transfer the thawed cell suspension to the conical tube containing 10 mL of growth medium.
3. Collect the cells by centrifugation at 1000 rpm for 5 minutes at room temperature. Remove the growth medium by aspiration.

4. Resuspend the cells in the conical tube in 15 mL of fresh growth medium by gently pipetting up and down.
5. Transfer the 15 mL of cell suspension to a T-75 tissue culture flask. Place the cells in a 37°C incubator at 5% CO₂.
6. Monitor cell density daily. Cells should be passaged when the culture reaches 95% confluence.

II. Freezing MEF Feeder Cells

1. Trypsinize cells and resuspend cell pellet in cold Freeze Medium at twice the desired final cell concentration.
2. Aliquot 1 mL of cells into sterile cryovials and place cryovials immediately into freezing container. Store overnight at -80°C.
3. Transfer frozen vials to -135°C freezer or liquid nitrogen.

III. Mitomycin C Treatment and Preparation of Feeder

1. Culture cells to 90% confluence. Wash it once with sterile PBS.
2. Add 10 µg/mL Mitomycin C (Sigma), incubate for 2 hrs.
3. Wash 3 times with sterile PBS to remove Mitomycin.
4. After dissociation by Trypsin, the Mitomycin-treated MEFs can be freezeed and stored in liquid nitrogen, or used as feeder by plating them at 75 000 cells/cm² in gelatin-coated tissue culture dishes for one day.

Recent Product Citation

Hai, Q. et al. (2022). Efficient Method to Differentiate Mouse Embryonic Stem Cells into Macrophages in vitro. *Bio-Protocol*. **12**(3):e4318. doi: 10.21769/BioProtoc.4318.

Warranty

These products are warranted to perform as described in their labeling and in Cell Biolabs literature when used in accordance with their instructions. THERE ARE NO WARRANTIES THAT EXTEND BEYOND THIS EXPRESSED WARRANTY AND CELL BIOLABS DISCLAIMS ANY IMPLIED WARRANTY OF MERCHANTABILITY OR WARRANTY OF FITNESS FOR PARTICULAR PURPOSE. CELL BIOLABS's sole obligation and purchaser's exclusive remedy for breach of this warranty shall be, at the option of CELL BIOLABS, to repair or replace the products. In no event shall CELL BIOLABS be liable for any proximate, incidental or consequential damages in connection with the products.

This product is for RESEARCH USE ONLY; not for use in diagnostic procedures.

Contact Information

Cell Biolabs, Inc.
7758 Arjons Drive
San Diego, CA 92126
Worldwide: +1 858-271-6500
USA Toll-Free: 1-888-CBL-0505
E-mail: tech@cellbiolabs.com
www.cellbiolabs.com

©2005-2022: Cell Biolabs, Inc. - All rights reserved. No part of these works may be reproduced in any form without permissions in writing.

