

Rabbit Anti-Caspase-9 Polyclonal Antibody

Cat # NB-30-00483

Description

Apoptosis is related to many diseases and induced by a family of cell death receptors and their ligands. Cell death signals are transduced by death domain containing adapter molecules and members of the caspase family of proteases. A novel member in the caspase family was recently identified and designated ICE-LAP6, Mch6, and Apaf-3. Caspase-9 and Apaf-1 bind to each other, which leads to caspase-9 activation. Caspase-9 is also activated by granzyme B and CPP32. Activated caspase-9 cleaves and activates caspase-3 that is one of the key proteases, being responsible for the proteolytic cleavage of many key proteins in apoptosis. Caspase-9 play a central role in cell death induced by a wide variety of apoptosis activators including TNFalpha, TRAIL, anti-CD-95, FADD, and TRADD. Caspase-9 is expressed in a variety of human tissues. The Cleaved Caspase-9 polyclonal antibody preferentially recognizes (active) cleaved Caspase-9. Whereas the antibody has a strong preference for (active) cleaved Caspase-9, in some cell or tissue systems or techniques the antibody may also recognize the pro form of Caspase-9.

Product informations

| | |
|--------------------------------|---|
| Clonality: | Polyclonal |
| Application: | WB, IHC, IP |
| Reactivity: | Dog, Gerbil, Human, Mouse, Rat |
| Gene: | CASP9 |
| Gene ID : | 842 |
| Uniprot ID : | P55211 |
| Format : | Sera |
| Alternative Name : | CASP9,MCH6 |
| Isotype : | Rabbit IgG |
| Immunogen Information : | A recombinant catalytically active of human Caspase-9 protein was used as the immunogen for this antibody |

Product Info

| | |
|---------------------------|--|
| Amount: | 50 µl |
| Content: | 50 µl sera |
| Storage condition: | Store the antibody at 4°C, stable for 6 months. For long-term storage, store at 20°C. Avoid repeated freeze and thaw cycles. |

Application Note

WB: 1:1000-1:2000, IHC (paraffin): 1:1000-1:5000, IHC (frozen): Users should optimize, IP: 1:50-1:200




Fig: 1 Formalin-fixed, paraffin-embedded dog brain sections stained for cleaved Caspase-9 expression using 20-1044 antibody at 1:2000. A and B. The pattern of Caspase-9 staining may vary between different types of neurons. Hematoxylin-eosin counterstain.




Fig:2 Formalin-fixed, paraffin-embedded dog brain sections stained for cleaved Caspase-9 expression using 20-1044 antibody at 1:2000. A. Section from a dog brain 2 hr after reperfusion injury. B. Section from a dog brain sham control (brain surgery but no injury). Hematoxylin-eosin counterstain.




Fig: 3 Formalin-fixed, paraffin-embedded human brain sections stained for cleaved Caspase-9 expression using 20-1044 antibody at 1:2000. A. Section from a patient 24 hr after head trauma. B. Control: section from a patient with no known neurological disease or head injury. Hematoxylin-eosin counterstain.