


Puromycin dihydrochloride [58-58-2]

#Cat: NB-64-11468-1mL	Size: 1mL
#Cat: NB-64-11468-10mg	Size: 10mg
#Cat: NB-64-11468-25mg	Size: 25mg
#Cat: NB-64-11468-50mg	Size: 50mg
#Cat: NB-64-11468-100mg	Size: 100mg



Chemical Properties

Cas No:	58-58-2
Formula:	C ₂₂ H ₃₁ Cl ₂ N ₇ O ₅
Molecular weight:	544.44
Appearance:	Solid
Storage:	store at low temperature Powder: -20°C for 3 years In solvent: -80°C for 1 year

Biological Description

Description	Puromycin dihydrochloride (CL13900 dihydrochloride) is a cinnamamide adenosine antibiotic and an inhibitor of protein synthesis. Puromycin dihydrochloride inhibits protein synthesis by binding to RNA and has antitumor and antitrypanosomal activity
Targets (IC₅₀)	ribosome, Antibacterial, Antibiotic, DNA/RNA Synthesis
In vitro	<p>METHODS: Human hepatocellular carcinoma cells, HepG2, and primary rat hepatocytes, PRH, were treated with Puromycin dihydrochloride (0-300 μmol/L) for 72 h, and cell growth inhibition was detected by MTT.</p> <p>RESULTS: Puromycin dihydrochloride dose-dependently inhibited the growth of HepG2 and PRH cells, with IC₅₀s of 1600 μmol/L and 2000 nmol/L, respectively.[1]</p> <p>METHODS: Human breast cancer cells MDA-MB-231 and MDA-MB-436 were transfected with shRNAs-Puro lentivirus, and after 24 h, the transfected cells were cultured in new medium containing Puromycin dihydrochloride (1 μg/ml) for about 12 days, and then the cells were screened for the success of transfection.</p> <p>RESULTS: Puromycin dihydrochloride screened the lentivirally transfected cells successfully. [2]</p> <p>METHODS: Cells transfected with lentivirus were cultured in medium supplemented with Puromycin dihydrochloride (2 μg/mL). RESULTS: Puromycin dihydrochloride screened and maintained lentivirally successfully transfected cells. [3]</p>
In vivo	Puromycin is utilized to select recombinant cells from non-cultured cells. It serves as a tool for studying protein synthesis in many systems.

Solubility Information

Solubility	H ₂ O: 92 mg/mL (168.98 mM), Sonication is recommended. Ethanol: < 1 mg/mL (insoluble or slightly soluble), DMSO: 255 mg/mL (468.37 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
-------------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.8367 mL	9.1837 mL	18.3675 mL
5 mM	0.3673 mL	1.8367 mL	3.6735 mL
10 mM	0.1837 mL	0.9184 mL	1.8367 mL
50 mM	0.0367 mL	0.1837 mL	0.3673 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Chan C, et al. Incompatibility of chemical protein synthesis inhibitors with accurate measurement of extended protein degradation rates. *Pharmacol Res Perspect*. 2017 Oct;5(5):e00359.

Wang Y N, Ruan D Y, Wang Z X, et al. Targeting the cholesterol-ROR α/γ axis inhibits colorectal cancer progression through degrading c-myc. *Oncogene*. 2022: 1-13.

Han W, et al. Roles of eIF3m in the tumorigenesis of triple negative breast cancer. *Cancer Cell Int*. 2020 Apr 29;20:141.

Gao C, Wang H, Wang T, et al. Platelet CLEC-2 regulates neuroinflammation and restores blood brain barrier integrity in a mouse model of traumatic brain injury. *Journal of Neurochemistry*. 2020: e14983

Wang YN, et al. Targeting the cholesterol-ROR α/γ axis inhibits colorectal cancer progression through degrading cmcyc. *Oncogene*. 2022 Dec;41(49):5266-5278.

Han W, Zhang C, Shi C, et al. Roles of eIF3m in the tumorigenesis of triple negative breast cancer. *Cancer Cell International*. 2020, 20(1): 1-16

Han W, Zhang C, Shi C, et al. Roles of eIF3m in the tumorigenesis of triple negative breast cancer[J]. *Cancer Cell International*. 2020, 20(1): 1-16.

Han W, Cao F, Gao X J, et al. ZIC1 acts a tumor suppressor in breast cancer by targeting survivin. *International Journal of Oncology*. 2018 Sep;53(3):937-948

Gao C, Wang H, Wang T, et al. Platelet CLEC-2 regulates neuroinflammation and restores blood brain barrier integrity in a mouse model of traumatic brain injury[J]. *Journal of Neurochemistry*. 2020: e14983.

Liu S, Zhang X, Yao X, et al. Mammalian IRE1 α dynamically and functionally coalesces with stress granules. *Nature Cell Biology*.2024: 1-15.

Li M, Lou L, Ren L, et al. EIF4G2 Promotes Hepatocellular Carcinoma Progression via IRES-dependent PLEKHA1 Translation Regulation.*Journal of Proteome Research*.2024

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use